

# СИБИРСКИЕ ЭЛЕКТРОННЫЕ МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 6, стр. 510–513 (2009)  
Краткие сообщения

УДК 514.8, 515.12  
MSC 28A80

## A SELF-SIMILAR CONTINUUM WHICH IS NOT THE ATTRACTOR OF ANY ZIPPER

A.V. TETENOV, O. PUREVDORJ

**ABSTRACT.** The article contains a construction of a self-similar dendrite which is not the attractor of any self-similar zipper.

**Keywords:** self-similar sets, fractal, self-similar dendrites, zippers.

### 1. INTRODUCTION

Let  $\mathcal{S}$  be a system  $\{S_1, \dots, S_m\}$  of injective contraction mappings of a complete metric space  $(X, d)$  to itself and let  $K$  be its *invariant set*, that is a non-empty compact set  $K$  satisfying  $K = \bigcup_{i=1}^m S_i(K)$ . The set  $K$  is also called the *attractor of the system*  $\mathcal{S}$ . There is a natural construction allowing to obtain the systems  $\mathcal{S}$  with an arcwise connected invariant set. This construction called a self-similar zipper goes back to the works of Thurston [5] and Astala [2] and was analyzed in detail by Aseev, Kravtchenko and Tetenov in [6]. Namely,

**Definition 1.** A system  $\mathcal{S} = \{S_1, \dots, S_m\}$  of injective contraction maps of complete metric space  $X$  to itself is called a zipper with vertices  $(z_0, \dots, z_m)$  and signature  $\vec{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_m) \in \{0, 1\}^m$  if for any  $j = 1, \dots, m$   $S_j(z_0) = z_{j-1+\varepsilon_j}$  and  $S_j(z_m) = z_{j-\varepsilon_j}$ .

If the maps  $S_i$  are similarities (or affine maps) the zipper is called self-similar (correspondingly self-affine).

We shall call the points  $z_0$  and  $z_m$  the *initial* and the *final* point of the zipper respectively.

TETENOV, A.V., PUREVDORJ, O, A SELF-SIMILAR CONTINUUM WHICH IS NOT THE ATTRACTOR OF ANY ZIPPER.

© 2009 TETENOV, A.V., PUREVDORJ, O.

The work is supported by RFFI (grant 09-01-90202).

Communicated by A.D. Mednykh November, 30, 2009, published December, 4, 2009.

The simplest example of a self-similar zipper may be obtained if we take a partition  $P$ ,  $0 = x_0 < x_1 < \dots < x_m = 1$  of the segment  $I = [0, 1]$  into  $m$  parts and put  $T_i = x_{i-1+\varepsilon_i}(1-t) + x_{i-\varepsilon_i}t$ . This zipper  $\{T_1, \dots, T_m\}$  will be denoted by  $\mathcal{S}_{P,\varepsilon}$ .

**Theorem 2.** (see [6]). *For any zipper  $\mathcal{S} = \{S_1, \dots, S_m\}$  with vertices  $\{z_0, \dots, z_m\}$  and signature  $\varepsilon'$  in a complete metric space  $(X, d)$  and for any partition  $0 = x_0 < x_1 < \dots < x_m = 1$  of the segment  $I = [0, 1]$  into  $m$  parts there exists a unique map  $\gamma : I \rightarrow K(\mathcal{S})$  such that for each  $i = 1, \dots, m$ ,  $\gamma(x_i) = z_i$  and  $S_i \cdot \gamma = \gamma \cdot T_i$  (where  $T_i \in \mathcal{S}_{P,\varepsilon}$ ), the map  $\gamma$  being Hölder continuous.*

The mapping  $\gamma$  in the Theorem 2 is called a *linear parametrization* of the zipper  $\mathcal{S}$ . Thus, the attractor  $K$  of any zipper  $\mathcal{S}$  is an arcwise connected set, whereas the linear parametrization  $\gamma$  may be viewed as a self-similar Peano curve, filling the continuum  $K$ .

**Some Peano curves.** The attractor  $K$  of a self-similar zipper  $\mathcal{S}$  with vertices  $(0, 0)$ ,  $(1/4, \sqrt{3}/4)$ ,  $(3/4, \sqrt{3}/4)$ ,  $(1, 0)$  and signature  $(1, 0, 1)$  is the Sierpinsky gasket.




FIGURE 1. Iterations 1,2,4, $\infty$  for Sierpinsky gasket.

A self-similar zipper with vertices  $(0, 0)$ ,  $(0, 1/2)$ ,  $(1/2, 1/2)$ ,  $(1, 1/2)$ ,  $(1, 0)$  and signature  $(1, 0, 0, 1)$  produces a self-similar Peano curve for the square  $[0, 1] \times [0, 1]$




FIGURE 2. Iterations 1,2,4 for square-filling Peano curve.

## 2. THE MAIN EXAMPLE.

The following example shows the existence of a self-similar continuum which cannot be represented as the attractor of a self-similar zipper.

Let  $\mathcal{S}$  be a system of contraction similarities  $g_k$  in  $\mathbb{R}^2$  where  $S_2(\vec{x}) = \vec{x}/2 + (2, 0)$ , and  $S_k(\vec{x}) = \vec{x}/4 + \vec{a}_k$  where  $\vec{a}_k$  run through the set  $\{(0, 0), (3, 0), (1, 2h), (3/2, 3h)\}$ ,  $h = \sqrt{3}/2$  for  $k = 1, 3, 4, 5$ . Let  $K$  be the invariant set of the system  $\mathcal{S}$  and  $T$  – the Hutchinson operator of the system  $\mathcal{S}$  defined by  $T(A) = \bigcup_{j=1}^5 S_j(A)$ .




FIGURE 3. Iterations 1,2,4,∞ for the example.

We shall use the following notation: By  $\Delta$  we denote the triangle with vertices  $A = (0, 0)$ ,  $B = (2, 2\sqrt{3})$  and  $C = (4, 0)$ , that is the convex hull of the points  $A, B$  and  $C$ . The point  $(2, 0)$  is denoted by  $D$ . Since for each  $S_i$ ,  $S_i(\Delta) \subset \Delta$ , the invariant set  $K$  of the system  $\mathcal{S}$  lies in  $\Delta$ . For a multiindex  $\mathbf{i} = i_1 \dots i_k$  we denote  $S_{\mathbf{i}} = S_{i_1} \dots S_{i_k}$ ,  $\Delta_{\mathbf{i}} = S_{\mathbf{i}}(\Delta)$ ,  $K_{\mathbf{i}} = S_{\mathbf{i}}(K)$ ,  $A_{\mathbf{i}} = S_{\mathbf{i}}(A)$ , etc.

1. *The set  $K$  is a dendrite.* The way the system  $\mathcal{S}$  is defined (see [3, Thm.1.6.2]) guarantees the arcwise connectedness of  $K$ . Since for each  $n$  the set  $T^n(\Delta)$  is simply-connected, the set  $K$  is a continuum, which contains no cycles, or a dendrite [4, Ch.6, §52]. Each point of  $K$  has the order 2 or 3. If a point  $x$  has the order 3, it is an image  $S_{\mathbf{i}}(D)$  of the point  $D$  for some multiindex  $\mathbf{i}$ . Any path in  $K$  connecting a point  $\xi \in J$  with a point  $\eta \in \Delta_{\mathbf{i}}$ ,  $\mathbf{i} = 4, 5, 24, 25, 224, 225, \dots$ , passes through the point  $D$ .

2. *Each non-degenerate segment  $J$ , contained in  $K$  is parallel to  $x$  axis and is contained in some maximal segment in  $K$  which has the length  $4^{1-n}$ .*

Consider a non-degenerate linear segment  $J \subset K$ . There is such multiindex  $\mathbf{i}$ , that  $J$  meets the boundary of  $S_{\mathbf{i}}(\Delta)$  in two different points which lie on different sides of  $S_{\mathbf{i}}(\Delta)$  and do not lie in the same subcopy of  $K_{\mathbf{i}}$ . Then  $J' = g_{\mathbf{i}}^{-1}(J \cap K_{\mathbf{i}})$  is a segment in  $K$  with endpoints lying on different sides of  $D$  which is not contained in neither of subcopies  $K_1, \dots, K_5$  of  $K$ . Then  $J' = [0, 4]$ . Since a part of  $J$  is a base of some triangle  $S_{\mathbf{i}}(\Delta)$ , the length of the maximal segment in  $K$  containing  $J$  is  $4^{1-n}$  where  $n \leq |\mathbf{i}|$ .

3. *Any injective affine mapping  $f$  of  $K$  to itself is one of the similarities  $S_{\mathbf{i}} = S_{i_1} \dots S_{i_k}$ .* Since  $f$  maps  $[0, 4]$  to some  $J \subset S_{\mathbf{i}}([0, 4])$  for some  $\mathbf{i}$ , it is of the form  $f(x, y) = (ax + b_1y + c_1, b_2y + c_2)$ , with positive  $b_2$ . Choosing appropriate composition  $S_{\mathbf{i}}^{-1} \cdot f \cdot S_{\mathbf{i}}(K)$  we obtain a map of  $K$  to itself sending  $[0, 4]$  to some subset of  $[0, 4]$ . Therefore we may suppose that  $f(x, y) = (ax + b_1y + c_1, b_2y)$ , and that the image  $f(\Delta)$  is contained in  $\Delta$  and is not contained in any  $\Delta_{\mathbf{i}}$ ,  $\mathbf{i} = 1, \dots, 5$ .

If  $f(B) \in \Delta_{\mathbf{i}}$ ,  $\mathbf{i} = 4, 5, 24, 25$ , then  $f(D) = D$  and  $c_1 = 2 - a$ .

If  $f(B) \in \Delta_i, i = 4, 5$ , then  $1/2 \leq b_2 \leq 1$ . In this case  $y$ -coordinates of the points  $f(B_1), f(B_3)$  are greater than  $\sqrt{3}/4$ , so they are contained in  $\Delta_1$  and  $\Delta_3$ , therefore the map  $f$  either keeps the points  $D_1, D_3$  invariant, or transposes them. In each case  $|a| = 1$  and  $f(\{A, C\}) = \{A, C\}$ . If in this case  $f(B) \neq B$ , then  $f(A_4)$  cannot be contained in  $T(\Delta)$ . The same argument shows that if  $f(B) = B$ , then  $f(A) \neq C$ . Therefore  $f = \text{Id}$ .

If  $f(B) \in \Delta_i, i = 24, 25$ , and  $a > 1/2$  then the points  $f(B_1), f(B_3)$  are again contained in  $\Delta_1$  and  $\Delta_3$ , therefore the map  $f$  either keeps the points  $D_1, D_3$  invariant, or transposes them, so  $|a| = 1$  and  $f(\{A, C\}) = \{A, C\}$ . Considering the intersections of the line segments  $[A, f(B)]$  and  $[f(B), C]$  with the boundary of  $T(\Delta)$  and  $T^2(\Delta)$  we see that either  $f(A_4)$  or  $f(C_5)$  are not contained in  $T^2(\Delta)$ , which is impossible.

Therefore, either  $a \leq 1/2$  or  $f = \text{Id}$ . The first means that  $f(\Delta) \subset \Delta_2$ , which contradicts the original assumption, so  $f = \text{Id}$ .

4. *The set  $K$  cannot be an attractor of a zipper.* Let  $\Sigma = \{\varphi_1, \dots, \varphi_m\}$  be a zipper whose invariant set is  $K$ . Let  $x_0, x_1$  be the initial and final points of the zipper  $\Sigma$ . Let  $\gamma$  be a path in  $K$  connecting  $x_0$  and  $x_1$ . Since for every  $i = 1, \dots, m$  the map  $\varphi_i$  is equal to some  $S_j$ , the sets  $\varphi_i(K)$  are the subcopies of  $K$ , therefore for each  $i$  at least one the images  $\varphi_i(x_0), \varphi_i(x_1)$  is contained in the intersection of  $\varphi_i(K)$  with adjacent copies of  $K$ . Consider the path  $\tilde{\gamma} = T_\Sigma(\gamma) = \bigcup_{i=1}^m \varphi_i(\gamma)$ . It starts from the point  $x_0$ , ends at  $x_1$  and passes through all copies  $K_j$  of  $K$ . Each of the points  $C_1 = A_2, C_2 = A_3, B_2 = C_4$  and  $B_4 = A_5$  splits  $K$  to two components, therefore is contained in  $\tilde{\gamma}$  and is a common point for the copies  $\varphi_i(\gamma), \varphi_{i+1}(\gamma)$  for some  $i$ . Therefore one of the points  $x_0, x_1$  must be  $A$ , one of the points  $x_0, x_1$  must be  $B$ , and one of the points  $x_0, x_1$  must be  $C$ , which is impossible.

## REFERENCES

- [1] Aseev V.V., *On the regularity of self-similar zippers*, Materials of the 6-th Russian-Korean Int. Symp. on Science and Technology, KORUS-2002 (June 24-30, 2002. Novosibirsk State Techn. Univ., Russia", Part 3, (Abstracts), 167.
- [2] Astala K., *Self-similar zippers*, Holomorph.funct. and Moduli: Proc.Workshop, March 13-19, 1986, New York, **1** (1988), 61–73.
- [3] Kigami J., *Analysis on fractals*, Cambridge University Press, 2001.
- [4] Kuratowski K., *Topology*, **2**, Academic Press, 1968.
- [5] Thurston W.P., *Zippers and univalent functions*, in: The Bieberbach Conjecture, Math. Surveys, No. 21, Am. Math. Soc., Providence, R.I. (1986), 185–197.
- [6] Aseev V.V., Kravtchenko A.S., Tetenov A.V., *On self-similar curves in the plane*, Siberian math J., **44**: 3 (2003), 481–492.
- [7] Hutchinson J., *Fractals and self-similarity*, Indiana Univ. Math. J., **30**: 5 (1981), 713–747.

ANDREY V. TETENOV,  
 GORNO-ALTAISK STATE UNIVERSITY,  
 LENKIN STR.,1, 649000, GORNO-ALTAISK, RUSSIA  
*E-mail address*: atet@mail.ru

OYUNAA PUREVDORJ,  
 KHOVD STATE UNIVERSITY,  
 KHOVD, MONGOLIA  
*E-mail address*: oyunaap@yahoo.com